

Problem 1

What is the upper 3 sigma confidence limit for proportion defective when the average daily production is 250 units with an established fraction defective of .05?

- a. .041
- b. .082
- c. .091
- d. .123

Problem 2

If the probability of a car starting on a cold morning is .6, and we have two such cars, what is the probability of at least one of the cars starting on a cold morning?

- a. .84
- b. .81
- c. .60
- d. .36

Problem 3

A sample of size 16 yields an average of 12 and a standard deviation of 3, estimate the 95% confidence interval for the population mean, μ .

- a. 10.4 to 13.6
- b. 10.45 to 13.55
- c. 10.53 to 13.47
- d. 10.77 to 13.23

Problem 4

Given the data following, what is the 90% confidence interval for the variance? [22, 23, 19, 17, 29, 25]

- a. 4.21 to 99.07
- b. 15.32 to 28.66
- c. 8.27 to 79.92
- d. 16.87 to 56.52

Problem 5

A process has a mean of 180 and a standard deviation of 8. What is the probability that a value selected at random will be between 170 and 200?

- a. .9937
- b. .8882
- c. .9937
- d. .8943

Problem 6

An experiment was conducted to compare two different tread designs. The data for tread life is shown below. Which test of hypothesis should be used first to compare the data?

- T test for paired comparisons
- T test assuming equal variances
- T test assuming unequal variances
- F test

Design A	Design B
44000	42000
43000	49000
47500	45500
44400	41800
46000	47300
45000	44750

Problem 7

Employee # Initial Mst Second Mst

1 78 81
2 73 76
3 66 60
4 45 60
5 89 90
6 59 75
7 78 82
8 87 92
9 69 66
10 91 94
11 55 61
12 77 77

Using the data above, can it be said at the 95% confidence level that the training improved the score?

- No, because t test is greater than t table
- No, because t test is less than t table
- Yes, because t test is greater than t table
- Yes, because t test is less than t table

Problem 8\

Given the ANOVA table shown we can conclude at the .01 level

- a. No significant difference between rows
- b. No significant difference between columns
- c. Significant difference between rows
- d. Significant difference between columns

<i>Source of Variation</i>	<i>SS</i>	<i>df</i>	<i>MS</i>	<i>F</i>	<i>P-value</i>	<i>F crit</i>
Rows	12.66667	4	3.166667	2.5	0.12572	7.0060
Columns	26.53333	2	13.26667	10.4736	0.00583	8.6491
Error	10.13333	8	1.266667			
Total	49.33333	14				

Problem 9

The slope regression equation for the following data is

X	Y
4	3
6	7
2	1
8	9
4	3
3	5
8	6
9	9
4	5
10	7
6	7

$$\Sigma X = 64$$

$$\Sigma Y = 62$$

$$\Sigma X^2 = 442$$

$$\Sigma XY = 416$$

- a. Not linear therefore no slope calculator
- b. 1.02
- c. .794
- d. .625